SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

MASAO NAGASAWA

Note on pasting of two Markov processes

Séminaire de probabilités (Strasbourg), tome 10 (1976), p. 532-535

<http://www.numdam.org/item?id=SPS_1976__10__532_0>

© Springer-Verlag, Berlin Heidelberg New York, 1976, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (http://portail.mathdoc.fr/SemProba/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

NOTE ON PASTING OF TWO MARKOV PROCESSES

By Masao NAGASAWA

The author remarked(e.g. in his lectures at UCSD in 1967/68 and at Erlangen in 1973/74) that Courrege-Priouret's theorem [1] of pasting two continuous strong Markov processes is a simple corollary of the theorem of piecing out(or revival) of a Markov process which was given by the author(cf.[3]). Since this remark was published nowhere, it would be of use to explain the procedure, though Meyer gave similar one recently in [4], which contains other interesting applications.

Let U and V be open subsets of a Polish space and adjoin ∂ as an extra point to U and V as usual. Suppose we have continuous strong Markov processes $(W^U, \underline{B}_t^U, x_t^U, P_x^U)$ on U and $(W^V, \underline{B}_t^V, x_t^V, P_x^V)$ on V which are killed at each boundary. Suppose $U \cap V \neq \phi$ and moreover

(1) x_t^U and x_t^V coincide on $U \cap V$ (the consistency condition),

that is, if we kill the two processes at the first leaving time T from $U \cap V$, we get the same Markov process on $U \cap V$. The problem is to construct a continuous strong Markov process x_t on $U \cup V$ pasting the two processes, in other wards, to obtain a process x_t which moves on x_t^U in U and on x_t^V in V. This pasting technique is an important tool, for example, when we construct diffusions on a Riemanian Manifold.

(i) Let $U \oplus V$ be the direct sum of U and V. When $x \in U \cap V$, denote x^U the corresponding point in U and x^V in V as a point in $U \oplus V$ (as is clear from Figure 1). The natural identification mapping p from $U \oplus V$ onto $U \cup V$ is defined by

$$p(x^U) = p(x^V) = x,$$

and we call it the pasting operator(not projection!).

Fig.1.



First we construct a process on $U \oplus V$ (treating U and V as separated sheets) and get the one on $U \cup V$ (pasting U and V). Let us define a continuous strong Markov process $(W^\circ, \underline{\underline{B}}^\circ_t, x^\circ_t, P^\circ_x)$ on $U \oplus V$, putting

$$W^\circ = W^U \oplus W^V, \qquad \underline{\underline{B}}^\circ_t = \underline{\underline{B}}^U_t \oplus \underline{\underline{B}}^V_t,$$

$$P^\circ_x = \begin{cases} P^U_x, & x \in U, \\ P^V_x, & x \in V, \end{cases} \qquad x^\circ_t = \begin{cases} x^U_t, & \text{on } W^U, \\ x^V_t, & \text{on } W^V. \end{cases}$$

Next, let us define a kernel on $(U \oplus V) \cup \{\partial\}$ by

$$n(x,\cdot) = \begin{cases} \varepsilon_{x^V} & \text{if } x = x^U \in (\partial U) \cap V, \\ \varepsilon_{x^U} & \text{if } x = x^V \in (\partial V) \cap U, \\ \varepsilon_\partial & \text{if otherwise,} \end{cases}$$

where ε_a denote the point mass at a, and put

$$N(w^\circ,\cdot) = \begin{cases} n(x^U_{\zeta-}(w^U),\cdot) & \text{if } w^\circ = w^U \in W^U, \\ n(x^V_{\zeta-}(w^V),\cdot) & \text{if } w^\circ = w^V \in W^V. \end{cases} \tag{2}$$

Then N is a "revival kernel" (it was called "instantaneous distribution" in [3]). Now apply the piecing out (or revival) theorem to x°_t and N, obtaining a right continuous strong Markov process $(\bar{W}, F_t, X_t, \bar{P}_x)$ on $U \oplus V$ (cf. figure 2). To obtain a continuous strong Markov process on $U \cup V$, that we are looking for, what we need to do is to apply the pasting operator p:

$$x_t = p(X_t), \qquad P_x = \bar{P}_{p^{-1}(x)}, \quad x \in U \cup V. \tag{3}$$



X_t on $U \oplus V$ Fig. 2. x_t on $U \cup V$

Since the pasting operator is a transformation of the state space (cf. [2], p.325), to be well-defined it is enough to prove

$$\bar{E}_{x^U}[f \circ p(X_t)] = \bar{E}_{x^V}[f \circ p(X_t)], \quad x \in U \cap V, \tag{4}$$

for f on $U \cup V$. The right hand side of (4) is equal to

$$\bar{E}_{x^U}[f \circ p(X_t); t<T] + \bar{E}_{x^U}[\bar{E}_{X_T}[f \circ p(X_{t-s})]|_{s=T}; t \geq T], \tag{5}$$

where T is the first leaving time from $U \cap V$. But from the way of construction of X_t, the first term of (5) is equal to

$$E_x^U[f(x_t^U); t<T],$$

which is equal to

$$E_x^V[f(x_t^V); t<T],$$

by the consistency condition (1), therefore is equal to

$$\bar{E}_{x_V}[f \circ p(X_t); t<T].$$

The second term of (5) is equal to

$$\bar{E}_{x_V}[\bar{E}_{X_T}[f \circ p(X_{t-s})]|_{s=T}; t \geqq T],$$

because

$$\bar{P}_{x_U}[X_T \in \cdot] = \bar{P}_{x_V}[X_T \in \cdot]. \tag{6}$$

Thus we have (4). It is clear that $x_t = p(X_t)$ is continuous.

(ii) Now, let x_t^U and x_t^V be strong Markov process on U and V, resp. We don't assume any continuity of paths, but assume $\exists\, x_{\zeta-}^U \in \bar{U}$ and $\exists\, x_{\zeta-}^V \in \bar{V}$, and the consistency condition (1). Let $n(x,dy)$ be a probability kernel on $\overline{(U \oplus V)} \cup \{\partial\}$ satisfying

$$\begin{aligned} &\text{the support of } n(x^U,\cdot) \text{ is contained in } (V \setminus U) \cup \{\partial\}, \\ &\text{the support of } n(x^V,\cdot) \text{ is contained in } (U \setminus V) \cup \{\partial\}. \end{aligned} \tag{7}$$

Define a revival kernel $N(w^\circ,\cdot)$ by (2) using this kernel $n(x,dy)$. To keep consistency for jumps of paths of the process which will be constructed, we assume the second consistency condition: When $x \in U \cap V$,

$$\begin{aligned} P_{x_U}^U[n(x_{\zeta-}^U,\cdot); T=\zeta] &= P_{x_V}^V[x_T^V \in \cdot; T<\zeta], \\ P_{x_V}^V[n(x_{\zeta-}^V,\cdot); T=\zeta] &= P_{x_U}^U[x_T^U \in \cdot; T<\zeta]. \end{aligned} \tag{8}$$

Let T_U and T_V be the first leaving times from U and V, resp., then

$x_t = p(X_t)$ is a strong Markov process on $U \cup V$ satisfying that x_t killed at T_U is equal to x_t^U, x_t killed at T_V is equal to x_t^V, and

$$\begin{aligned} P_z[x_{T_U} \in \cdot \,|\, x_{T_U-} = x] &= n(x,\cdot), \quad \text{if } x \in U, \\ P_z[x_{T_V} \in \cdot \,|\, x_{T_V-} = x] &= n(x,\cdot), \quad \text{if } x \in V. \end{aligned} \tag{9}$$

If x_t^U and x_t^V are right continuous, x_t on $U \cup V$ is also right continuous. The equality (4) is proved in the same way. The equality (6) in this

case is verified as follows: Putting $a = x^U$ and $b = x^V$, $x \in U \cap V$,

$$\begin{aligned}
\bar{P}_a[X_T \in \cdot] &= \bar{P}_a[X_T \in \cdot; T=r_1] + \bar{P}_a[X_T \in \cdot; T<r_1] \\
&= P_x^U[n(x^U_{\zeta-}, \cdot); T=\zeta] + P_x^U[x^U_T \in \cdot; T<\zeta] \\
&= P_x^V[x^V_T \in \cdot; T<\zeta] + P_x^V[n(x^V_{\zeta-}, \cdot); T=\zeta] \\
&= \bar{P}_b[X_T \in \cdot; T<r_1] + \bar{P}_b[X_T \in \cdot; T=r_1] \\
&= \bar{P}_b[X_T \in \cdot],
\end{aligned}$$

where r_1 is the first jumping(or revival) time of X_t. The second and the fourth equalities are by the revival theorem, and the third is by the second consistency condition (8). (9) is clear because

$$P_z[x_{r_1} \in \cdot | x_{T_U-}=x] = \bar{P}_z[X_{r_1} \in \cdot | X_{r_1-}=x] = n(x,\cdot).$$

(iii) Appendix.

DEFINITION. A probability kernel $N(w,dy)$ on $W \times \underline{\underline{B}}(S)$ is revival kernel (or instantaneous distribution) if it satisfies

(i) $N(\theta_t w,\cdot) = N(w,\cdot)$, if $t < \zeta(w)$, and

(ii) $N(w,\cdot) = \varepsilon_\partial$, if $\zeta(w) = 0$.

THEOREM (of revival or piecing out). Let $(W, \underline{\underline{B}}_t, x_t, P_x)$ $x \in S$ be a Markov process and N be a revival kernel. Then there exists a Markov process $(\bar{W}, \underline{\underline{F}}_t, X_t, \bar{P}_x)$ on S satisfying

(i) X_t, $t < r_1$, is equivalent to x_t,

(ii) $\bar{P}_x[X_{r_1} \in \cdot | \underline{\underline{F}}_{r_1-}] = N(w_1,\cdot)$,

where $\bar{W} = W^{(1,2,3,\cdots)} \ni \bar{w} = (w_1,w_2,\cdots)$ and $r_1(\bar{w}) = \zeta(w_1)$. If x_t is strong Markov , so is X_t, and if x_t is right continuous, so is X_t. If x_t is continuous and if $N(w,\cdot) = \varepsilon_{x_{\zeta-}(w)}$, then X_t is continuous.

References.

[1] Courrège,Ph. et P.Priouret; Recollements de processus de Markov. Publ.Inst.Statist.Univ. Paris 14(1965)275-377.

[2] Dynkin,E.B.; Markov processes, Springer-Verlag 1965.

[3] Ikeda,N. M.Nagasawa, and S.Watanabe; Branching Markov processes II Journal Math. Kyoto Univ. 8(1968)365-410.

[4] Meyer P.A.; Renaissance, recollements, melanges, ralentissement de processus de Markov. Annales de l'Inst. Fourier, to appear.