SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

GARETH C. PRICE
DAVID WILLIAMS

Rolling with "slipping" : I

Séminaire de probabilités (Strasbourg), tome 17 (1983), p. 194-197
<http://www.numdam.org/item?id=SPS_1983__17__194_0>

© Springer-Verlag, Berlin Heidelberg New York, 1983, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (http://portail.mathdoc.fr/SemProba/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM
Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
http://www.numdam.org/

ROLLING WITH 'SLIPPING' : I

by

Gareth C. Price and David Williams

Throughout this note, ∂ will denote the Stratonovich differential, and d will denote the Itô differential.

Let B be a $BM(\mathbb{R}^3)$, a Brownian motion on $\mathbb{R}^3$. Let Z be a process on $\mathbb{R}^3$ with $|Z(0)| = 1$ and

$$\partial Z = Z \times \partial B, \tag{1}$$

where $\times$ is the vector product. Then Z is a $BM(S^2)$, a Brownian motion on the unit sphere S^2 in $\mathbb{R}^3$.

Though the representation (1) of a $BM(S^2)$ is very simple, it appears to suffer from the disadvantage that "there is too much freedom in B". Only the component

$$\partial Y = \partial B - (Z.\partial B)Z$$

of ∂B 'tangential' to S^2 really matters. Now, of course, we have

$$\partial Y = (\partial Z) \times Z$$

and the proper driving equation for Z:

$$\partial Z = Z \times \partial Y.$$

The apparent defect of redundancy in (1) is illusory. We have the very satisfying situation that the equation

$$\partial b = Z.\partial B, \qquad b(0) = 0,$$

defines a $BM(\mathbb{R}^1)$ process b <u>independent</u> of the process Z. We think of b as providing the information about B which is missing from Z.

Suppose now that $\tilde{Z}$ is a $BM(S^2)$ adapted to the filtration of Z. Let

$$\partial \tilde{Y} = (\partial \tilde{Z}) \times \tilde{Z}.$$

THEOREM. <u>We have</u>

$$d\tilde{Y} = H\,dY \qquad \text{(an <u>Itô</u> equation)},$$

where

(i) for each t, H_t is an orthogonal transformation such that

$$H_t Z_t = \tilde{Z}_t,$$

(ii) the process $H = \{H_t\}$ is Z previsible.

Proof that Z is a $BM(S^2)$. We have $\partial(Z.Z) = 2Z.\partial Z = 0$, so that Z stays on S^2. It is clear that Z is invariant under $O(3)$, whence Dynkin's formula shows that the infinitesimal generator of Z is a constant multiple of the spherical Laplacian. Of course, the usual argument for Stratonovich equations shows that if G is the generator of Z, then, with $\partial_k = \partial/\partial Z_k$,

$$2G = (Z_3\partial_2 - Z_2\partial_3)^2 + (Z_1\partial_3 - Z_3\partial_1)^2 + (Z_2\partial_1 - Z_1\partial_2)^2,$$

and this 'squared angular momentum' operator is known to be the Laplacian on S^2. □

Proof that b is a $BM(\mathbb{R}^1)$ independent of Z. The generator of the 4-dimensional process (b,Z) is

$$\tfrac{1}{2}(Z_1\partial_b + Z_3\partial_2 - Z_2\partial_3)^2 + \text{two cyclic permutations},$$

where $\partial_b = \partial/\partial b$. This operator splits as

$$\tfrac{1}{2}\partial_b^2 + G.$$ □

Martingale characterization of $BM(S^2)$. A process U on $\mathbb{R}^3$ with $|U(0)| = 1$ is a $BM(S^2)$ if and only if U is a continuous semimartingale such that

(i) $dU + Udt = dM$, where M is a martingale,

(ii) $d\langle U_m, U_n \rangle = (\delta_{mn} - U_m U_n)dt$.

A 'trick' way to prove this is to apply stereographic projection of S^2 to $\mathbb{R}^2 \cup \{\infty\}$, and then apply the Stroock-Varadhan result to the resulting process on $\mathbb{R}^2$.

Proof of Theorem. By Jacod's Theorem that "martingale characterization implies martingale representation" (see [1], and especially result (11) there), we have

$$d\tilde{Z}_m + \tilde{Z}_m dt = A^{(m)}.dN \tag{2}$$

where $A^{(m)}$ is a Z previsible process with values in $\mathbb{R}^3$, and

$$dN = dZ + Zdt = Z \times dB. \tag{3}$$

Thus,

$$d\tilde{Z}_m + \tilde{Z}_m dt = K^{(m)}.dB,$$

where

$$K^{(m)} = A^{(m)} \times Z.$$

We have

$$d\left\langle \tilde{Z}_m, \tilde{Z}_n \right\rangle /dt = \delta_{mn} - \tilde{Z}_m \tilde{Z}_n = K^{(m)}.K^{(n)},$$

and, also, the vectors $K^{(m)}$ are perpendicular to Z. But, the vectors

$$J^{(1)} = (0, -\tilde{Z}_3, \tilde{Z}_2),\ J^{(2)} = (\tilde{Z}_3, 0, -\tilde{Z}_1),\ J^{(3)} = (-\tilde{Z}_2, \tilde{Z}_1, 0) \tag{4}$$

satisfy

$$J^{(m)}.J^{(n)} = \delta_{mn} - \tilde{Z}_m \tilde{Z}_n,$$

and the vectors $J^{(m)}$ are perpendicular to $\tilde{Z}$. Hence, for some (in fact, unique) orthogonal matrix H_t with $H_t Z_t = \tilde{Z}_t$,

$$K_t^{(m)} = J_t^{(m)} H_t.$$

The process H is Z previsible, and

$$d\tilde{B} = HdB$$

defines a B Brownian motion. On combining (2), (3), and (4), we find that

$$d\tilde{Z} + \tilde{Z}dt = \tilde{Z} \times d\tilde{B}, \quad \partial\tilde{Z} = \tilde{Z} \times \partial\tilde{B}.$$

Of course, since H is orthogonal,

$$\tilde{Z}.d\tilde{B} = Z.dB,$$

so that, with an obvious notation, we have the satisfying relation

$$\tilde{b} = b.$$

Next,

$$\partial\tilde{Y} = (\partial\tilde{Z}) \times \tilde{Z}, \qquad d\tilde{Y} = (d\tilde{Z}) \times \tilde{Z},$$

so that

$$d\tilde{Y} = (\tilde{Z} \times d\tilde{B}) \times \tilde{Z} = d\tilde{B} - (\tilde{Z}.d\tilde{B})\tilde{Z}$$

$$= d\tilde{B} - (Z.dB)\tilde{Z} = HdY,$$

and our theorem is proved. □

The title indicates the 'frame-bundle' significance, something we hope to explain in a wider context. There seem to be some very nice - and potentially important - applications.

Acknowledgement. We wish to thank Professor A. Truman for helpful discussions on these topics. A.T. and D.W. intend to publish follow-up work. [The representation

$$dX = n(X) \times dB$$

for Brownian motion on a surface ($n(\cdot)$ is the unit normal) allows some neat formulae.]

REFERENCE

[1] J. JACOD, A general theorem of representation for martingales, Probability (ed. J.L. Doob), Proc. Symp. Pure Math. Amer. Math. Soc. XXXI (1977), 37-54.

Department of Mathematics
University College
SWANSEA SA2 8PP
Great Britain